

Orogenic climax of Earth: The 1.2–1.1 Ga Grenvillian superevent

Martin J. Van Kranendonk^{1,2} and Christopher L. Kirkland^{3,4}

¹School of Biological, Earth and Environmental Sciences, and Australian Centre for Astrobiology, University of New South Wales, Kensington, NSW 2052, Australia

²ARC Centre of Excellence for Core to Crust Fluid Systems; Administering Institution, Department of Earth and Planetary Sciences, Macquarie University, NSW 2109, Australia

³Geological Survey of Western Australia, 100 Plain Street, East Perth, WA 6004, Australia

⁴Department of Imaging and Applied Physics, Curtin University, Perth, WA 6845, Australia

ABSTRACT

The rate of growth of the continental crust is controversial. We present an evaluation of time-constrained analyses of oxygen isotopes in zircon grains and incompatible element (Zr, Th) concentrations in magmatic rocks to test for variations in the degree of crustal recycling through geological time. The data indicate a rise in these geochemical proxies from ca. 3.0 Ga to a statistically significant peak at 1.2–1.1 Ga during the amalgamation of supercontinent Rodinia, and a decrease thereafter. When combined with other geological and geophysical observations, the data are interpreted as a consequence of an unprecedented level of crustal recycling and sediment subduction during Rodinia assembly, arising from a “Goldilocks” (i.e., just right) combination of larger, thicker plates on a warmer Earth with more rapid continental drift relative to modern Earth. The subsequent decrease in $\delta^{18}\text{O}$, Zr, and Th measurements is interpreted to reflect decreasing drift rates on a cooling Earth.

INTRODUCTION

The growth of Earth's continental crust over time depends on the rate of production at arcs and in oceanic plateaus versus the rate of recycling back into the mantle through subduction erosion. Current estimates suggest that these competing processes have been approximately in balance since the Neoproterozoic (Scholl and von Heune, 2009), implying that continental crust reached a maximum volume at some time in the Precambrian, when the Earth was hotter.

However, the rates of crustal growth over geological time remain controversial. Armstrong (1991) advocated for the essentially complete growth of continental crust prior to 3.5 Ga, whereas others have suggested more gradual crustal growth (e.g., Belousova et al., 2010). Furthermore, it is unknown whether crust grew in pulses associated with periods of supercontinent assembly (Stein and Hofmann, 1994; Condie, 1998), or whether the episodic record is an artifact of preservation (e.g., Hawkesworth et al., 2009). Nonetheless, recent studies indicating periods of enhanced exchange between the lower and upper mantle lend support to the episodic nature of crustal growth (Condie et al., 2001; Roberts, 2012; Van Kranendonk et al., 2012).

Incompatible trace elements (ITEs) represent another way of investigating the growth and/or recycling of crust, as they are preferentially excluded from common rock-forming minerals during fractional crystallization due to their unsuitable size and/or charge to fit within the cation sites of crystals, and are partitioned into silicate liquids during melting. This process has led to average upper continental crust highly enriched in ITEs relative to the mantle (Zr = 237 vs. 11 ppm, Th = 11 vs. 0.09 ppm). Elevated concentrations of Zr and Th above these crustal values in magmas reflect enhanced crustal recycling and the importance of preexisting continental crust in the orogenic cycle.

Oxygen isotope variations also reflect the orogenic cycle through the nature of the source material: low values ($\delta^{18}\text{O} = 5.3\text{‰} \pm 0.3\text{‰}$) imply a mantle source, or a crustal source not exposed to surface weathering, whereas higher values (7.5‰–12‰) imply incorporation of material affected by near-surface weathering processes into magmas through subduction (Valley et al., 2005).

Naturally, these proxies will vary spatially at any one time, but global data sets should identify temporal changes in source contributions, as well as potential changes in the amount of crustal recycling, as ITEs are preferentially concentrated and oxygen isotopes elevated toward heavy values by subduction, collision-related uplift, erosion, and crustal recycling.

In this paper we investigate the changing nature of the global supercontinent cycle by analysis of a global compilation of $\delta^{18}\text{O}$ values from dated zircon grains, complemented by ITE concentrations (Zr and Th) in a global database of dated magmatic rocks (Appendices DR1 and DR2 in the GSA Data Repository¹). The data show statistically significant peaks in oxygen isotope and ITE values at 1.2–1.1 Ga, accompanying the Grenvillian assembly of supercontinent Rodinia, and a successive decrease thereafter. We argue that the data reflect a global climax in the amount of crustal recycling arising from a “Goldilocks” (i.e., just right) combination of fast drift of larger, and possibly thicker, continental plates on a warmer Earth with more rapid continental drift compared with modern day.

METHODS AND SAMPLE MATERIALS

We present whole-rock concentrations of Zr and Th from 340 samples of magmatic rocks of a range of ages from Western Australia; the magmatic crystallization ages were determined by U-Pb analysis of zircon through secondary ionization mass spectrometry (Appendix DR1). These data are complemented by a global compilation of 2913 analyses of Zr and Th concentrations in magmatic rocks, extracted from the Petrological Database (PetDB; <http://www.earthchem.org/petdb>), North American Volcanic and Intrusive Rock Database (NAVDAT; <http://www.navdat.org>), and the Geochemistry of Rocks of the Oceans and Continents database (GEOROC; <http://georoc.mpch-mainz.gwdg.de/georoc>) (as queried through the Integrated Earth Data Applications EarthChem website; <http://www.earthchem.org>). Oxygen isotope data are compiled from 1262 dated magmatic zircon grains from previously published sources (Appendix DR2).

To analyze the data, we employ the scan statistic (Naus, 1965; Glaz et al., 2009) to retrospectively search through time for data clusters with high values, using a normal probability model (Kulldorff, 1997; Kulldorff et al., 2009) in which a temporal window of all sizes between 5 and 100 Ma is moved over the time series. The maximum value relative to the number of observations in the window is recorded over all possible time periods and compared to its distribution under the null hypothesis of a purely random process (for more detail, see the Data Repository).

RESULTS

When plotted against time, the $\delta^{18}\text{O}$, Zr, and Th data sets show low-amplitude peaks and troughs after ca. 3.0 Ga that correlate with periods of supercontinent assembly and dispersal as determined from geological

¹GSA Data Repository item 2013205, Appendix DR1 (whole rock chemical parameters and SIMS U-Pb zircon crystallization age for magmatic rocks within Western Australia), and Appendix DR2 (oxygen isotope values in zircon grains from the compilation of Valley et al. [2005] with additional data from Kirkland et al. [2010] and Be'eri-Shlevina et al. [2009]), is available online at www.geosociety.org/pubs/ft2013.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.




Figure 1. Evolution plots. A: $\delta^{18}\text{O}$ in zircon grains from global compilation (as detailed in Appendix DR2; see footnote 1) set against backdrop of histogram of zircon ages (from Condie, 1998). VSMOW—Vienna standard mean ocean water. **B, C:** Concentrations of incompatible trace elements (Zr and Th) in dated igneous rocks from global geochemical data sets (<http://www.earthchem.org>) augmented by dated rocks from Western Australia (Appendix DR1; and see West Australian Resource Information and Map Services Geochemistry Database Search: <http://www.dmp.wa.gov.au/geochem>). Supercontinents: P—Pangea; G—Gondwana; R—Rodinia; N—Nuna; S/S—Superia/Sclavia.

and isotopic data (Fig. 1). Using the scan statistic, a statistically significant ($p < 0.001$) temporal cluster of heavy oxygen and high Zr concentration is present at 1.2–1.1 Ga, during Grenvillian assembly of the supercontinent Rodinia (Table DR1 in the Data Repository). Scanning across Proterozoic Earth history (2.5–0.5 Ga) reveals a statistically significant ($p < 0.001$) temporal cluster of high Th concentration, also at 1.2–1.1 Ga.

These peaks are not only higher at 1.2–1.1 Ga than at any time during the preceding 2.5 Ga, but also higher than at any time thereafter (Fig. 1). Oxygen isotope concentrations in zircon were on average 1.84% greater than at any other time in Earth history; whole-rock Zr and Th concentrations were on average 332 ppm and 22 ppm greater than at any other time in all of Earth history, and in Proterozoic Earth history, respectively (Table DR1). The 1.2–1.1 Ga peak identified herein is also evident in Zr concentrations from the Grenville orogen of North America (to 1700–1800 ppm; Moecher and Samson, 2006).




Figure 2. Schematic evolution of Grenvillian supercontinent. A: Leading to Rodinia assembly, showing multiple subduction zones (green lines, teeth on hanging wall) and collisions (red areas). **B:** Final Rodinia assembly across Grenville orogen. Widespread subduction during supercontinent assembly results in mantle cooling.

DISCUSSION

Orogenic Climax in Earth History

The 1.2–1.1 Ga peak in geochemical and isotopic signatures, identified in Western Australia, North America, and global data sets, indicates that the Grenvillian orogeny represents a unique episode in Earth evolution. Enrichment of ITEs may arise from fractionation and/or multiple episodes of remelting of crustal sources. In the case of Zr, temperature is an important control on Zr solubility and its concentration in melts (Moecher and Samson, 2006). However, these processes can only proceed so far and generally only produce maximum concentrations of 500–600 ppm Zr in A-type granites, with rare maxima of 1370 ppm (Collins et al., 1982; Whalen et al., 1987), well below the maximum values presented here and in Moecher and Samson (2006). It is significant that melting temperature is regarded as having no control on the variation of oxygen isotopes through time (Valley et al., 2005). Combined, these considerations imply a distinct orogenic control on the compositional anomalies identified here that relate to the total volume of subducted sediment and/or recycled continental crust.

Five aspects of the Grenvillian orogenic cycle are anomalous and have direct bearing on the geochemical and oxygen isotopic peaks identified here.

1. Large tectonic plates previously formed during dispersal of the supercontinent Nuna ca. 1.8 Ga were made significantly larger by addition of juvenile crust over wide areas from 1.7 to 1.3 Ga, within marginal plate settings of arcs and backarcs (e.g., southwest and southeast North America, Baltica, southwest Australia; Dickinson and Higgins, 1992; Slagstad et al., 2009; Evans and Mitchell, 2011; Kirkland et al., 2011).

2. Large areas of the continental crust were also substantially thickened over the same time interval by underplating basaltic magmas that fractionated to generate voluminous, temporally unique, Mesoproterozoic anorthosite-mangerite complexes (Windley, 1995).

3. Earth's mantle was $\sim 55^\circ\text{C}$ hotter than present (Labrosse and Jaupart, 2007; Michaut and Jaupart, 2007; Brown, 2008), as evidenced by the lack of blueschists in orogens older than the Neoproterozoic and the presence of thicker ophiolites in the Mesoproterozoic (Moore, 1993).

4. The rate of continental drift immediately prior to Grenvillian orogenesis was nearly an order of magnitude larger than at any subsequent time in Earth history (O'Neill et al., 2007; Swanson-Hysell et al., 2009).

Although primarily a function of hotter more rapidly convecting mantle, higher drift rates at this time may have been exacerbated by enhanced slab pull of thicker oceanic lithosphere (i.e., Moores, 1993).

5. Paleogeographic reconstructions and tectonic analysis reveal that the Grenville orogen was perhaps the longest and widest in Earth history (giant-supergiant orogen of Beaumont et al., 2010), spanning 25% of the globe, or a distance of ~20,000 km along strike and as wide as 800 km, including a core zone several hundred kilometers wide with large slabs of deep-crustal eclogite (Fig. 2; Eaton et al., 1995; Rivers et al., 2002; Li et al., 2008). These features indicate a vast area of crustal thickness doubling during a protracted continent-continent collision (Davidson, 2008; Hynes and Rivers, 2010). For comparison, the largest and highest orogen on modern Earth is the Alpine-Himalayan orogen, which is ~8000 km long and as much as 400 km wide.

These features demonstrate that the Grenville orogen was of unprecedented size, necessitating a powerful driving force. Whereas Beaumont et al. (2010) ascribed variations in orogenic magnitude solely to variations in temperature, data from the Himalaya orogen show that it is the product of the anomalously rapid drift of the Indian continent (18–20 cm/yr; Kumar et al., 2007), dragged on top of a large subducting plate of cold oceanic lithosphere. Thus, continental drift rate is an additional, and perhaps highly significant, factor in determining orogenic magnitude, with temperature likely to be a resultant function of rollback-generated backarc extension (Collins, 2002). Plate size and strength (rigidity) also play a significant role in orogen magnitude: small, weak (hot) plates produce small, or broad, low-grade orogens, whereas large, stiff plates produce large orogens (Tapponnier et al., 1982; Cruden et al., 2006). Combined, rapid drift rate, enhanced plate size and strength, and elevated temperature define what we refer to here as high orogenic intensity.

The long duration and large aerial extent of juvenile crust formation in arc-backarc settings immediately preceding Grenvillian orogenesis, combined with the evidence for multiple collisions during orogeny (1.3–1.0 Ga; Davidson, 2008; Hynes and Rivers, 2010), are important in understanding the peak in geochemical and isotopic anomalies recorded here (Fig. 2). Large-scale extension and rollback of the subducting plates creates juvenile crust in backarcs, whereas pulses of compression during accretion episodes consume newly generated crust (Gower and Tucker, 1994; Fisher et al., 2010). Such tectonic switching generates large volumes of ITE-enriched sediment derived from the continental hinterland and erosion of newly formed arc and backarc crust in developing mountain belts (Curry, 1994; Collins, 2002). This sediment, affected by high degrees of surficial weathering and transported into subduction zones, becomes incorporated into the magmatic system and fractionated (Chiarenzelli et al., 2010).

It is this tectonic switching, operating at the unprecedented scale of the Grenville orogen and in combination with the fast drift rate, anomalously thick and more rigid continental plates, and thicker oceanic lithosphere involved in Rodinia assembly, that we suggest was the cause for the unrivalled level of recycled orogenic crust at this time. That the scale of this process was evidently unmatched throughout the rest of Earth history suggests that this period represents a “Goldilocks” combination of large plates on a warmer, more rapidly convecting Earth (Fig. DR1).

Prelude

Secular cooling of Earth since at least the Neoproterozoic (Herzberg et al., 2010) has contributed to the changing style and thermal gradients of more recent orogenic belts (Brown, 2008). Such temporal changes in tectonic style are the result of large plates on cooler mantle, which have different rheological properties, styles of interaction, and insulating capacities than smaller plates on hotter mantle (Anderson, 1982; Rey and Houseman, 2006; Hynes and Rivers, 2010). Through the Archean, the combination of higher overall geothermal gradients and higher radiogenic heat production within continental crust resulted in weak, generally

submergent continental plates and wide, hot orogens (Cruden et al., 2006) that were unable to produce the scale of orogeny, or associated crustal recycling as on younger Earth.

By 2.4 Ga continents had emerged (Rey and Coltice, 2008), by 2.0 Ga bona fide ophiolites appeared in the rock record (e.g., Kontinen, 1987), and the modern-style supercontinent Nuna had formed by 1.8 Ga (Reddy and Evans, 2009); all of this points to the progressive thickening and stiffening of continental lithosphere through the Paleoproterozoic. However, Nuna assembly did not produce the same degree of crustal recycling shown by Rodinia, implying either that pre-Nuna plates were not as large as, and/or weaker than, post-Nuna plates, or drift rates were slower. Given the evidence of a nearly global magmatic shutdown at 2.45–2.2 Ga (Condie et al., 2009), the latter may have been the most significant factor (Fig. DR1).

Dénouement

The decrease in geochemical and isotopic proxies of orogenic recycling after the Grenvillian superevent is here ascribed to secular decrease in mantle temperatures and resultant slower rates of continental drift and thinner oceanic lithosphere. In addition, recent estimates of tectonic erosion at subduction zones exceeding crustal growth at arcs suggest that continental plates may have actually decreased in volume since 1.0 Ga (Scholl and von Heune, 2009), thereby further contributing to reduced orogenic intensity after this period.

CONCLUSIONS

A statistically significant ($p < 0.01$) and unrivalled cluster of high values in geochemical proxies of crustal recycling (zircon $\delta^{18}\text{O}$, whole-rock Zr and Th) is found at 1.2–1.1 Ga, representing an orogenic climax in Earth history during the Grenvillian assembly of supercontinent Rodinia. We suggest that this climax reflects a “Goldilocks” combination of large plates on a warmer mantle with faster rates of continental drift. Prior to that time, plates were smaller and less rigid and thus unable to match the orogenic intensity of the Grenvillian superevent. Post-Grenvillian secular decrease in mantle heat, slower drift rates, and thinner oceanic lithosphere, coupled with an imbalance of subduction erosion over crust production, resulted in smaller continental plates and declining orogenic intensity after 1.2–1.1 Ga.

ACKNOWLEDGMENTS

Analytical work was carried out under the auspices of the Government of Western Australia. M. Prause drafted the figures. R. Smithies, W. Collins, B. Murphy, K. Condie, and P. Hoffman provided critical reviews that helped clarify the manuscript. This is contribution 22 from the ARC Centre of Excellence for Core to Crust Fluid Systems. Kirkland publishes with permission of the executive director of the Geological Survey of Western Australia.

REFERENCES CITED

- Anderson, D.L., 1982, Hotspots, polar wander, Mesozoic convection and the geoid: *Nature*, v. 297, p. 391–393, doi:10.1038/297391a0.
- Armstrong, R.L., 1991, The persistent myth of continental growth: *Australian Journal of Earth Sciences*, v. 38, p. 613–630, doi:10.1080/08120099108727995.
- Beaumont, C., Jamieson, R., and Nguyen, M., 2010, Models of large, hot orogens containing a collage of reworked and accreted terranes: *Canadian Journal of Earth Sciences*, v. 47, p. 485–515, doi:10.1139/E10-002.
- Belousova, E.A., Kostitsyn, Y.A., Griffin, W.L., Begg, G.C., O'Reilly, S.Y., and Pearson, N.J., 2010, The growth of continental crust: Constraints from Hf-isotope data: *Lithos*, v. 119, p. 457–466, doi:10.1016/j.lithos.2010.07.024.
- Brown, M., 2008, Characteristic thermal regimes of plate tectonics and their metamorphic imprint throughout Earth history: When did Earth first adopt a plate tectonics mode of behavior?, *in* Condie, K.C., and Pease, V., eds., When did plate tectonics begin on planet Earth?: Geological Society of America Special Paper 440, p. 97–128, doi:10.1130/2008.2440(05).
- Chiarenzelli, J., Lupulescu, M., Cousens, B., Thern, E., Coffin, L., and Regan, S., 2010, Enriched Grenvillian lithospheric mantle as a consequence of long-lived subduction beneath Laurentia: *Geology*, v. 38, p. 151–154, doi:10.1130/G30342.1.
- Collins, W.J., 2002, Hot orogens, tectonic switching, and creation of continental crust: *Geology*, v. 30, p. 535–538, doi:10.1130/0091-7613(2002)030<0535:HOTSAC>2.0.CO;2.

- Collins, W.J., Beams, S.D., White, A.J.R., and Chappell, B.W., 1982, Nature and origin of A-type granites with particular reference to southeastern Australia: *Contributions to Mineralogy and Petrology*, v. 80, p. 189–200, doi:10.1007/BF00374895.
- Condie, K.C., 1998, Episodic continental growth and supercontinents: A mantle avalanche connection?: *Earth and Planetary Science Letters*, v. 163, p. 97–108, doi:10.1016/S0012-821X(98)00178-2.
- Condie, K.C., Des Marais, D.J., and Abbott, D.H., 2001, Precambrian superplumes and supercontinents: A record in black shales, carbon isotopes, and paleoclimates?: *Precambrian Research*, v. 106, p. 239–260, doi:10.1016/S0301-9268(00)00097-8.
- Condie, K.C., O'Neill, C., and Aster, R.C., 2009, Evidence and implications for a widespread magmatic shutdown for 250 My on Earth: *Earth and Planetary Science Letters*, v. 282, p. 294–298, doi:10.1016/j.epsl.2009.03.033.
- Cruden, A.R., Nasser, M., and Pysklywec, R., 2006, Surface topography and internal strain variation in wide hot orogens from three-dimensional analogue and two-dimensional numerical vice models, *in* Buitter, S.J.H., and Schreurs, G., eds., *Analogue and numerical modelling of crustal-scale processes: Geological Society of London Special Publication 253*, p. 79–104, doi:10.1144/GSL.SP.2006.253.01.04.
- Curry, J.R., 1994, Sediment volume and mass beneath the Bay of Bengal: *Earth and Planetary Science Letters*, v. 125, p. 371–383, doi:10.1016/0012-821X(94)90227-5.
- Davidson, A., 2008, Late Paleoproterozoic to mid-Neoproterozoic history of northern Laurentia: An overview of central Rodinia: *Precambrian Research*, v. 160, p. 5–22, doi:10.1016/j.precamres.2007.04.023.
- Dickin, A.P., and Higgins, M.H., 1992, Sm/Nd evidence for a major 1.5 Ga crust-forming event in the Central Grenville Province: *Geology*, v. 20, p. 137–140, doi:10.1130/0091-7613(1992)020<0137:SNEFAM>2.3.CO;2.
- Eaton, D.W., Hynes, A., Indares, A., and Rivers, T., 1995, Seismic images of eclogites, crustal-scale extension, and Moho relief in the eastern Grenville province, Quebec: *Geology*, v. 23, p. 855–858, doi:10.1130/0091-7613(1995)023<0855:SIOECS>2.3.CO;2.
- Evans, D.A.D., and Mitchell, R.N., 2011, Assembly and breakup of the core of Paleoproterozoic–Mesoproterozoic supercontinent Nuna: *Geology*, v. 39, p. 443–446, doi:10.1130/G31654.1.
- Fisher, C.M., Loewy, S.L., Miller, C.F., Berquist, P., Van Schmus, W.R., Hatcher, R.D. Jr., Wooden, J.L., and Fullagar, P.D., 2010, Whole-rock Pb and Sm–Nd isotopic constraints on the growth of southeastern Laurentia during Grenvillian orogenesis: *Geological Society of America Bulletin*, v. 122, p. 1646–1659, doi:10.1130/B30116.1.
- Glaz, J., Pozdnyakov, V., and Wallenstein, S., 2009, Scan statistics: Methods and applications: New York, Springer, 394 p.
- Gower, C.F., and Tucker, R.D., 1994, Distribution of pre-1400 Ma crust in the Grenville province: Implications for rifting in Laurentia–Baltica during Geon 14: *Geology*, v. 22, p. 827–830, doi:10.1130/0091-7613(1994)022<0827:DOPMCI>2.3.CO;2.
- Hawkesworth, C.J., Cawood, P.A., Kemp, A.I.S., Storey, C., and Dhuime, B., 2009, A matter of preservation: *Science*, v. 323, p. 49–50, doi:10.1126/science.1168549.
- Herzberg, C., Condie, K., and Korenaga, J., 2010, Thermal history of the Earth and its petrological expression: *Earth and Planetary Science Letters*, v. 292, p. 79–88, doi:10.1016/j.epsl.2010.01.022.
- Hynes, A., and Rivers, T., 2010, Protracted continental collision—Evidence from the Grenville orogen: *Canadian Journal of Earth Sciences*, v. 47, p. 591–620, doi:10.1139/E10-003.
- Kirkland, C.L., Spaggiari, C.V., Pawley, M.J., Wingate, M.T.D., Smithies, R.H., Howard, H.M., Tyler, I.M., Belousova, E.A., and Poujol, M., 2011, On the edge: U–Pb, Lu–Hf, and Sm–Nd data suggests reworking of the Yilgarn craton margin during formation of the Albany–Fraser Orogen: *Precambrian Research*, v. 187, p. 223–247, doi:10.1016/j.precamres.2011.03.002.
- Kontinen, A.T., 1987, An Early Proterozoic ophiolite—The Jormua mafic-ultramafic complex, northern Finland: *Precambrian Research*, v. 35, p. 313–341, doi:10.1016/0301-9268(87)90061-1.
- Kulldorff, M., 1997, A spatial scan statistic: *Communications in Statistics Theory and Methods*, v. 26, p. 1481–1496, doi:10.1080/03610929708831995.
- Kulldorff, M., Huang, L., and Konty, K., 2009, A scan statistic for continuous data based on the normal probability model: *International Journal of Health Geographics*, v. 8, p. 58, doi:10.1186/1476-072X-8-58.
- Kumar, P., Yuan, X., Kumar, M.R., Rainer Kind, R., Li, X., and Chadha, R.K., 2007, The rapid drift of the Indian tectonic plate: *Nature*, v. 449, p. 894–897, doi:10.1038/nature06214.
- Labrosse, S., and Jaupart, C., 2007, Thermal evolution of the earth: Secular changes and fluctuations of plate characteristics: *Earth and Planetary Science Letters*, v. 260, p. 465–481, doi:10.1016/j.epsl.2007.05.046.
- Li, Z.X., and 17 others, 2008, Assembly, configuration, and break-up history of Rodinia: A synthesis: *Precambrian Research*, v. 160, p. 179–210, doi:10.1016/j.precamres.2007.04.021.
- Michaut, C., and Jaupart, C., 2007, Secular cooling and thermal structure of continental lithosphere: *Earth and Planetary Science Letters*, v. 257, p. 83–96, doi:10.1016/j.epsl.2007.02.019.
- Moecher, D.P., and Samson, S.D., 2006, Differential zircon fertility of source terranes and natural bias in the detrital zircon record: Implications for sedimentary provenance analysis: *Earth and Planetary Science Letters*, v. 247, p. 252–266, doi:10.1016/j.epsl.2006.04.035.
- Moore, E.M., 1993, Neoproterozoic oceanic crustal thinning, emergence of continents, and origin of the Phanerozoic ecosystem: A model: *Geology*, v. 21, p. 5–8, doi:10.1130/0091-7613(1993)021<0005:NOCTEO>2.3.CO;2.
- Naus, J.I., 1965, The distribution of the size of the maximum cluster of points on the line: *American Statistical Association Journal*, v. 60, p. 532–538, doi:10.1080/01621459.1965.10480810.
- O'Neill, C., Lenardic, A., Moresi, L., Torsvik, T.H., and Lee, C.-T.A., 2007, Episodic Precambrian subduction: *Earth and Planetary Science Letters*, v. 262, p. 552–562, doi:10.1016/j.epsl.2007.04.056.
- Reddy, S.M., and Evans, D.A.D., 2009, Palaeoproterozoic supercontinents and global evolution: Correlations from core to atmosphere, *in* Reddy, S.M., et al., eds., *Palaeoproterozoic supercontinents and global evolution: Geological Society of London Special Publication 323*, p. 1–26, doi:10.1144/SP323.1.
- Rey, P.F., and Coltice, N., 2008, Neoproterozoic lithosphere strengthening and the coupling of Earth's geochemical reservoirs: *Geology*, v. 36, p. 635–638, doi:10.1130/G25031A.1.
- Rey, P.F., and Houseman, G., 2006, Lithospheric scale gravitational flow: The impact of body forces on orogenic processes from Archaean to Phanerozoic, *in* Buitter, S.J.H., and Schreurs, G., eds., *Analogue and numerical modelling of crustal-scale processes: Geological Society of London Special Publication 253*, p. 153–167, doi:10.1144/GSL.SP.2006.253.01.08.
- Rivers, T., Ketchum, J., Indares, A., and Hynes, A., 2002, The high pressure belt in the Grenville province: Architecture, timing, and exhumation: *Canadian Journal of Earth Sciences*, v. 39, p. 867–893, doi:10.1139/e02-025.
- Roberts, N.M.W., 2012, Increased loss of continental crust during supercontinent amalgamation: *Gondwana Research*, v. 21, p. 994–1000, doi:10.1016/j.gr.2011.08.001.
- Scholl, D.W., and von Heune, R., 2009, Implications of estimated magmatic additions and recycling losses at the subduction zones of accretionary (non-collisional) and collisional (suturing) orogens, *in* Cawood, P.A., and Kröner, A., eds., *Earth accretionary systems in space and time: Geological Society of London Special Publication 318*, p. 105–125, doi:10.1144/SP318.4.
- Slagstad, T., Culshaw, N.G., Daly, J.S., and Jamieson, R.A., 2009, Western Grenville province holds key to midcontinental Granite-Rhyolite Province enigma: *Terra Nova*, v. 21, p. 181–187, doi:10.1111/j.1365-3121.2009.00871.x.
- Stein, M., and Hofmann, A.W., 1994, Mantle plumes and episodic crustal growth: *Nature*, v. 372, p. 63–68, doi:10.1038/372063a0.
- Swanson-Hysell, N.L., Maloof, A.C., Weiss, B.P., and Evans, D.A.D., 2009, No asymmetry in geomagnetic reversals recorded by 1.1-billion-year-old Keweenaw basalts: *Nature Geoscience*, v. 2, p. 713–717, doi:10.1038/ngeo622.
- Tapponnier, P., Peltzer, G., Le Dain, A.Y., Armijo, R., and Cobbold, P., 1982, Propagating extrusion tectonics in Asia: New insights from simple experiments with plasticine: *Geology*, v. 10, p. 611–616, doi:10.1130/0091-7613(1982)10<611:PETIAN>2.0.CO;2.
- Valley, J.W., and 12 others, 2005, 4.4 billion years of crustal maturation: Oxygen isotope ratios of magmatic zircon: *Contributions to Mineralogy and Petrology*, v. 150, p. 561–580, doi:10.1007/s00410-005-0025-8.
- Van Kranendonk, M.J., Altermann, W., Beard, B.L., Hoffman, P.F., Johnson, C.J., Kasting, J.F., Melezhik, V.A., Nutman, A.P., Papineau, D., and Pirajno, F., 2012, A chronostratigraphic division of the Precambrian: Possibilities and challenges, *in* Gradstein, F.M., et al., eds., *The geologic time scale 2012: Amsterdam, Elsevier*, p. 299–392, doi:10.1016/B978-0-444-59425-9.00016-0.
- Whalen, J.B., Currie, K.L., and Chappell, B.W., 1987, A-type granites: Geochemical characteristics, discrimination and petrogenesis: *Contributions to Mineralogy and Petrology*, v. 95, p. 407–419, doi:10.1007/BF00402202.
- Windley, B.F., 1995, *The evolving continents* (third edition): New York, John Wiley and Sons, 526 p.

Manuscript received 18 November 2012

Revised manuscript received 23 January 2013

Manuscript accepted 3 February 2013

Printed in USA